Saskatchewan
Ministry of
Agriculture

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 20, August 18, 2011

For the Period August 9 to 15, 2011

Harvest is underway according to Saskatchewan Agriculture's weekly Crop Report. Three per cent of the 2011 crop has been combined. The five-year (2006-2010) provincial average for this time of year is five per cent combined.

One year ago
Less than one per cent of the 2010 crop had been combined. Six per cent of the winter wheat and fall rye and three per cent of the lentils and peas had been combined.

Twenty-one per cent of the winter wheat, 30 per cent of the fall rye, 11 per cent of the lentils and 10 per cent of the field peas have been combined. Twelve per cent of the canola, six per cent of the mustard, and one per cent of the spring wheat, oats and barley has been swathed.

Livestock producers in many areas of the province are finished haying, however rainfall delayed progress in some areas this past week. The average provincial hay yield for both dryland alfalfa and alfalfa/brome stands is 1.9 tons per acre, which is the five-year average (2006 to 2010). Average yield on irrigated alfalfa and alfalfa/brome stands is 2.9 tons per acre. Average greenfeed yield is 2.2 tons per acre on dryland and 3.4 tons per acre on irrigated stands.

Thunderstorms throughout the province brought heavy rain, high wind, and damaging hail in some areas. The majority of crop damage is due to hail, wind and insects.

Across the province, topsoil moisture on cropland is rated as eight per cent surplus, 82 per cent adequate and 10 per cent short. Hay land and pasture topsoil moisture is rated as four per cent surplus, 80 per cent adequate and 16 per cent short.

Farmers are busy swathing, combining and hauling hay.

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Tel: (306) 694-3592. Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

Thunderstorms rolled through the region with varying amounts of precipitation recorded. Precipitation ranged from nil to 35 mm for the week. High winds also caused some swathed crops to blow. The Frobisher area received 20 mm, the Stoughton and Marquis areas 6 mm, the Moosomin area 30 mm, the Tantallon area 21 mm, the Grenfell area 5 mm, the Weyburn area 35 mm, the Pangman area 25 mm, the Pense area 21 mm, the Radville area 25 mm and the Ceylon area 22 mm. According to the Crop Report precipitation information, the Moosomin area has the third highest cumulative rainfall since April 1 at 470 mm. This falls shortly behind the Bengough area at 484 mm and the Dorintosh area in the northwest which is reporting 547 mm.

Although rain moved through the region, some producers were able to get back onto the combines. Three per cent of the 2011 crop has been harvested. Ten per cent is swathed or ready to straight combine. Eleven per cent of the lentils, 15 per cent of the peas, 51 per cent of the winter wheat, 40 per cent of the fall rye and two per cent of the canola have been combined. Twenty-one per cent of the canola and eight per cent of the mustard has been swathed. Crop conditions and staging vary throughout the region, depending on the spring's excess moisture and seeding date.

Many crop reporters are indicating it is difficult to assess yields at this time on those crops that have not been combined, due to varying crop stages, flooded areas and disease. The following average yields for the region are predicted as follows: winter wheat 45 bu/ac, spring wheat 33 bu/ac, durum 29 bu/ac, oats 52 bu/ac, barley 45 bu/ac, fall rye 42 bu/ac, flax 14 bu/ac, canola 26 bu/ac, mustard 800 lbs/ac, lentils 1000 lbs/ac, peas 29 bu/ac and canaryseed 1100 lbs/ac.

Crop reporters have indicated that hay yields are variable across the region. Estimated average hay yields are as follows: dry land alfalfa and alfalfa/brome stands 2.1 tons per acre, dry land greenfeed 1.9 tons per acre and other tame hay 1.8 tons per acre.

Topsoil moisture rating on cropland is reported as 11 per cent surplus, 77 per cent adequate and 12 per cent short. On hay land and pasture, topsoil moisture is rated as three per cent surplus, 78 per cent adequate and 19 per cent short. Crop district 3ASE is reporting 45 and 31 per cent short of topsoil moisture on crop land and hay and pasture, respectively.

The majority of crop damage is due to wind, hail and insects. High winds blew some canola and hay swaths around. Hail damage was reported in the Moosomin and Marquis areas. Diamondback moth and Bertha Armyworm have also caused crop damage in canola in some areas. Crops in some of the areas have been damaged by the hot temperatures experienced in July and into August. Crops planted under excess moisture conditions developed shallow root systems that were not able to buffer against high temperatures.

Producers are busy finishing haying, hauling bales, getting ready for harvest, combining, swathing and controlling weeds on unseeded acres.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

Thunderstorms rolled through some areas in the region dropping large amounts of rain, slowing harvest operations. Precipitation reported ranged from 3 to 84 mm. Most areas in crop district 3ASW reported rain over 35 mm. The Bengough area reported 44 mm, the Rockglen area 46 mm, the Glentworth area 84 mm, the Lafleche area 17 mm, the Chaplin area 30 mm, the Hazenmore area 39 mm, the Ponteix and Shaunavon areas 26 mm, the Swift Current area 10 mm, the Eastend area 12 mm and the Fox Valley area 24 mm. According to our crop report rainfall database, the Bengough area has the second highest cumulative rainfall (since April 1) for the province at 484 mm (The Dorintosh area in the north west has 547 mm total cumulative rainfall since April 1).

Seven per cent of the 2011 crop has been combined. Thirteen per cent has been swathed or is ready to straight cut. Twenty per cent of the lentils, 22 per cent of the field peas, 19 per cent of the winter wheat, four per cent of the canola and one per cent of the mustard have been combined.

Many crop reporters are indicating it is difficult to assess yields at this time on those crops that have not been combined, due to varying crop stages and wet areas. The following average yields for the region are as follows: winter wheat 35 bu/ac, spring wheat 30 bu/ac, durum 32 bu/ac, oats 51 bu/ac, barley 47 bu/ac, fall rye 39 bu/ac, flax 17 bu/ac, canola 25 bu/ac, mustard 1000 lbs/ac, lentils 1300 lbs/ac, peas 28 bu/ac, canaryseed 1000 lbs/ac and chickpeas 1300 lbs/ac.

Estimated average hay yields are predicted as follows: dry land alfalfa 1.9 tons per acre, alfalfa/brome stands 2.1 tons per acre, dry land greenfeed 2.2 tons per acre and other tame hay 1.8 tons per acre, irrigated alfalfa and alfalfa/brome 3 tons per acre.

Cropland topsoil moisture is rated as three per cent surplus, 77 per cent adequate, 19 per cent short and one per cent very short. Hay land and pasture topsoil moisture conditions are 72 per cent adequate and 28 per cent short. Crop district 4A is reporting 35 and 40 per cent short on topsoil moisture in crop land and hay and pasture land, respectively.

Hail and high winds caused the majority of the crop damage. Many areas reported hail in crop districts 3ASW and 3BS. Quarter sized hail fell near the Glentworth area for about 30 minutes significantly damaging crops in the area. Crop reporters have indicated hail damage in other areas of the region ranges from minimal to 100 per cent.

The recent rain will help fill some of the later seeded crops. Producers are busy finishing haying, harvesting pulses and winter cereals, swathing, and controlling weeds in chemfallow and summerfallow.

East-Central Saskatchewan (Crop Districts 5 and 6a)

Thunderstorms were reported and most areas received rain again this week, however amounts range from nil to 77 mm. High winds and hail were also reported causing swaths to blow around and damage to crops. Crop districts 5A and 5B received more rain than crop district 6A. The Stockholm area received 44 mm, the Langenburg area 60 mm, the Saltcoats area 75 mm, the Jedburgh area 18 mm, the Ituna area 45 mm, the Raymore area 7 mm, the Elfros area 46 mm, the Kelvington area 34 mm, the Bethune area 19 mm, the Craik area 5 mm and the Jansen area 6 mm.

Less than one per cent of the 2011 crop has been combined in the region. Ten per cent has been swathed or is ready to straight cut. Five per cent of the lentils, three per cent of the peas, 12 per cent of the winter wheat and 24 per cent of the fall rye have been combined. Fourteen per cent of the canola and five per cent of the mustard has been swathed.

Many crop reporters are indicating it is difficult to assess yields at this time. The following average yields for the region are predicted as follows: winter wheat 48 bu/ac, spring wheat 36 bu/ac, durum 34 bu/ac, oats 76 bu/ac, barley 56 bu/ac, fall rye 41 bu/ac, flax 23 bu/ac, canola 30 bu/ac, mustard 1100 lbs/ac, lentils 1100 lbs/ac, peas 37 bu/ac and canaryseed 1000 lbs/ac.

Estimated average hay yields are as follows: dry land alfalfa and alfalfa/brome 1.9 tons per acre, dry land greenfeed 2.4 tons per acre and other tame hay 1.9 tons per acre.

Topsoil moisture on cropland is reported as 15 per cent surplus, 75 per cent adequate and 10 per cent short. Hay land and pasture topsoil moisture is rated as 14 per cent surplus, 71 per cent adequate and 15 per cent short.

Crop damage from high winds and hail was reported in many areas. Bertha armyworm in canola is also causing crop damage. The heavy rain that fell in some areas was not needed as producers start their harvest operations. Hail damage on crops varied from very little to 75 per cent. Crops in crop district 6A are further advanced than those in 5A and 5B.

Farmers are busy finishing haying, hauling bales, swathing and desiccating pulses and combining.

West-Central Saskatchewan (Crop Districts 6B and 7)

Spotty thunderstorms slowed haying and harvest operations for a little while in some areas. Precipitation reported ranged from nil to 58 mm. The Langham area recorded 32 mm, the Sonningdale area 15 mm, the Rosetown and Kindersley areas 17 mm, the Harris area 31 mm, the Smiley area 29 mm, the Biggar area 58 mm, the Landia area 23 mm, the Scott area 13 mm.

Less than one per cent of the 2011 crop has been harvested. Three per cent has been swathed or is ready to straight cut. Two per cent of the lentils, peas and winter wheat have been combined, 17 per cent of the fall rye has been combined. Six per cent of the canola has been swathed.

Many crop reporters are indicating it is difficult to assess yields at this time. The following average yields for the region are predicted as follows: winter wheat 43 bu/ac, spring wheat 36 bu/ac, durum 35 bu/ac, oats 70 bu/ac, barley 56 bu/ac, fall rye 38 bu/ac, flax 25 bu/ac, canola 33 bu/ac, mustard 1200 lbs/ac, lentils 1300 lbs/ac, peas 38 bu/ac and canaryseed 1200 lbs/ac.

Estimated average hay yields are as follows: dry land alfalfa 2 tons per acre, alfalfa/brome 1.8 tons per acre, dry land greenfeed 2.2 tons per acre and other tame hay 1.5 tons per acre, irrigated alfalfa 2.4 tons per acre and irrigated alfalfa/brome 2.5 tons per acre, irrigated greenfeed 3.3 tons per acre.

Topsoil moisture on cropland is reported as four per cent surplus, 89 per cent adequate and seven per cent short. Hay land and pasture topsoil moisture is rated as four per cent surplus, 82 per cent adequate and 14 per cent short.

Many areas in the region reported hail and wind damage to crops. Hail damage up to 100 per cent loss has been reported, other areas reported minor crop damage. Bertha armyworm is also causing crop damage in canola.

Farmers are busy finishing haying, hauling bales, desiccating pulse crops, swathing and getting ready for harvest.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

The north east region recorded some rain over the past week, but not as much as areas to the south. Most areas reported rain in amounts ranging from nil to 14 mm. The Tisdale area reported 14 mm, the Codette area 7 mm, the Nipawin and Prince Albert areas 3 mm, the Alvena area 10 mm and the Melfort area 6 mm.

A few fields of winter cereals have been combined. Less than one per cent of the 2011 crop has been swathed or is ready to straight cut. Less than one per cent of the canola has been swathed.

Many crop reporters are indicating it is difficult to assess yields at this time. The following average yields for the region are predicted as follows: winter wheat 45 bu/ac, spring wheat 42 bu/ac, oats 82 bu/ac, barley 66 bu/ac, fall rye 32 bu/ac, flax 23 bu/ac, canola 34 bu/ac and peas 36 bu/ac.

Estimated average hay yields are as follows: dry land alfalfa 2.1 tons per acre, alfalfa/brome 2 tons per acre, dry land greenfeed 3.4 tons per acre and other tame hay 1.7 tons per acre.

Cropland topsoil moisture is rated as six per cent surplus, 90 per cent adequate and four per cent short. Hay land and pasture topsoil moisture is rated as four per cent surplus, 87 per cent adequate, seven per cent short and two per cent very short.

Of the crop damage reported, the majority is due to high winds that have lodged some crops and moved some canola swaths around.

Farmers are busy finishing haying, hauling bales, swathing and getting ready for harvest.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

A week of very little rain is just what producers needed to continue with haying operations in the northwest region. Most areas recorded amounts below 4 mm. The Duck Lake and Dorintosh areas were the only two areas that recorded into the double digits at 12 and 32 mm, respectively. The Radisson and North Battleford areas received 4 mm, the Meota area 5 mm and the Frenchman Butte area 7 mm. According to the Crop Report rainfall information, the area around Dorintosh has the highest amount of cumulative rainfall in the province since April 1, at 547 mm.

A few fields of winter cereals have been combined. Two per cent of the canola has been swathed and two per cent of the pea crops have been desiccated and are ready to straight cut.

Many crop reporters are indicating it is difficult to assess yields at this time. The following average yields for the region are predicted as follows: winter wheat 40 bu/ac, spring wheat 41 bu/ac, oats 86 bu/ac, barley 70 bu/ac, fall rye 33 bu/ac, flax 20 bu/ac, canola 35 bu/ac and peas 41 bu/ac.

Estimated average hay yields are as follows: dry land alfalfa 1.4 tons per acre, alfalfa/brome 1.3 tons per acre, dry land greenfeed 1.8 tons per acre and other tame hay 1.1 tons per acre.

Topsoil moisture conditions for cropland are four per cent surplus, 94 per cent adequate and two per cent short. Pasture and hay land conditions are two per cent surplus, 94 per cent adequate, three per cent short and one per cent very short.

The majority of crop damage is due to insects and disease. Grasshoppers, bertha armyworm are causing crop damage.

Although crops are behind normal in development, the warm weather has significantly helped advance maturity. Producers are busy haying, monitoring fields for insects and getting ready for harvest.

Saskatchewan Harvest Progress August 15, 2011				
winter wheat	**% Standing**	**% in swath**	**% ready to straight combine**	**% combined**
southeast	30	1	18	51
southwest	50	2	30	18
east central	53	8	27	12
west central	90	0	8	2
northeast	N/A	N/A	N/A	N/A
northwest	N/A	N/A	N/A	N/A

Saskatchewan Harvest Progress August 15, 2011				
Fall rye	**% Standing**	**% in swath**	**% ready to straight combine**	**% combined**
southeast	25	31	5	39
southwest*	38	10	6	34
east central	21	10	45	24
west central**	46	35	0	17
northeast	N/A	N/A	N/A	N/A
northwest	N/A	N/A	N/A	N/A
* 12 per cent is rated as 'other' **2 per cent is rated as 'other'				

Saskatchewan Harvest Progress August 15, 2011				
Lentils	**% Standing**	**% in swath**	**% ready to straight combine**	**% combined**
southeast	73	9	7	11
southwest*	41	15	22	20
east central	77	2	16	5
west central	93	1	4	2
northeast	100	0	0	0
northwest	N/A	N/A	N/A	N/A
*2 per cent rated as 'other'				

Saskatchewan Harvest Progress August 15, 2011				
Peas	**% Standing**	**% in swath**	**% ready to straight combine**	**% combined**
southeast*	61	7	16	15
southwest	54	6	18	22
east central	79	8	10	3
west central	91	1	6	2
northeast	94	2	4	0
northwest	98	0	2	0
* one per cent rated as 'other'				

Saskatchewan Harvest Progress August 15, 2011				
Canola	**% Standing**	**% in swath**	**% ready to straight combine**	**% combined**
southeast	77	21	0	2
southwest	61	35	0	4
east central	86	14	0	0
west central	94	6	0	0
northeast	99	1	0	0
northwest	99	1	0	0

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending August 15, 2011



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2011 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 17, 2011

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period August 9 to 15, 2011

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	N/A	240
	2	Mount Pleasant	N/A	294
	3	Enniskillen	20	334
	33	Moose Creek	N/A	291
	34	Browning	16	384
	61	Antler	N/A	305
	63	Moose Mountain	15	301
	64	Brock	N/A	272
	85	Tecumseh	6	424
1B	91	Maryfield	6	385
	92	Walpole	N/A	282
	95	Golden West	18	342
	122	Martin	30	470
	123	Silverwood	16	357
	124 A	Kingsley	4	309
	124 B	Kingsley	N/A	N/A
	125	Chester	N/A	287
	151 A	Rocanville	21	443
	151 B	Rocanville	N/A	346
	153	Willowdale	N/A	347
	154	Elcapo	N/A	286
	156	Wolseley	5	348
2A	66	Griffin	N/A	316
	67	Weyburn	21	464
	68	Brokenshell	35	453
	69	Norton	25	386
	97	Wallington	26	411
2B	127 A	Francis	1	342
	127 B	Francis	2	259
	129	Bratt's Lake	1	260
	156	Indian Head	NIL	336
	160 A	Pense	N/A	171
	160 B	Pense	20.5	236
	161	Moose Jaw	4	170
	191	Marquis	6	195
3ASE	7	Souris Valley	N/A	N/A
	36 A	Laurier	14	394
	36 B	Laurier	25	433
	39 A	The Gap	3	314
	39 B	The Gap	22	402
3ASW	10	Happy Valley	28	485
	11	Hart Butte	N/A	83
	12	Poplar Valley	N/A	310
	40	Bengough	44	484
	42	Willow Bunch	36	394
	43	Old Post	46	384
	44 A	Waverley	37	292
	44 B	Waverley	84	311
	70	Key West	N/A	346
	71	Excel	N/A	374
	73 A	Stonehenge	6.09	335
	73 B	Stonehenge	N/A	364
	74	Wood River	16.5	349
3AN	17	Val Marie	75	494
	18	Lone Tree	N/A	259
	45	Maple Creek	34	466
	75 A	Pinto Creek	39	337
	75 B	Pinto Creek	24	347
	76 A	Auvergne	18	256
	76 B	Auvergne	26	262
	77	Wise Creek	N/A	270
	78 A	Grassy Creek	12	317
	78 B	Grassy Creek	13	316
	78 C	Grassy Creek	26	269
	101	Terrell	4	272
	102	Lake Johnston	7.5	291
	103	Sutton	N/A	320
	105	Glenbain	22	332
	106	Whiska Creek	11	275
	107	Lac Pelletier	N/A	230
	108	Bone Creek	12	307
	134	Shamrock	N/A	270
	164	Chaplin	30	302
	193 A	Eyebrow	NIL	203
	193 B	Eyebrow	NIL	204
	194	Enfield	N/A	219
3DN	138	Webb	8	208
	166	Excelsior	N/A	241
	167	Sask. Landing	3	295
	168	Riverside	10	200
	226	Victory	N/A	198
	228	Lacadena	8	196
	255	Coteau	N/A	N/A
	257	Monet	5	169

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	109	Carmichael	8	284
	110	Piapot	N/A	139
	111	Maple Creek	N/A	143
	49 A	White Valley	12	209
	49 B	White Valley	5	201
	51	Reno	3.2	315
	79	Arlington	2	227
4B	139	Gull Lake	6	185
	141	Big Stick	9	167
	142	Enterprise	6	190
	169	Pittville	N/A	153
	171	Fox Valley	24	175
	229	Miry Creek	N/A	N/A
	231	Happyland	12	202
5A	183	Fertile Belt	44	456
	186	Abernethy	16	273
	211	Churchbridge	60	367
	213	Saltcoats	75	356
	216	Tullymet	30	270
	217	Lipton	14	217
	241	Calder	77	306
	243	Wallace	2	255
	245 A	Garry	16	236
	245 B	Garry	7	351
	245 C	Garry	8	319
	246	Ituna Bon Accord	45	281
	247	Kellross	22	268
	248	Touchwood	7	180
5B	271	Cote	3	197
	276	Foam Lake	N/A	42
	277	Emerald	NIL	262
	305	Invermay	2	285
	307	Elfros	46	356
	308	Big Quill	2	181
	333 A	Clayton	N/A	NIL
	333 B	Clayton	N/A	240
	334	Preeceville	N/A	213
	335	Hazel Dell	12	269
	336	Seaman	22	281
	337	Lakeview	13	282
	338	Lakeside	7	207
	366	Kelvington	34	288
6A	189	Lumsden	N/A	219
	190 A	Dufferin	19	247
	190 B	Dufferin	2	233
	190 C	Dufferin	15	180
	190 D	Dufferin	N/A	197
	219	Longlaketon	N/A	175
	220	Mckillop	1	194
	221	Sarnia	0.4	233
	222	Craik	1	220
	251	Big Arm	3	211
	252	Arm River	5	183
	279	Mount Hope	N/A	231
	280	Wreford	3	224
	282	Mccraney	4	238
	309	Prairie Rose	6	208
	313	Lost River	N/A	139
	339	Leroy	0.2	189
	340	Wolverine	1	178
	341	Viscount	N/A	205
	343	Blucher	5	169
6B	283	Rosedale	2	128
	284	Rudy	NIL	243
	285	Fertile Valley	NIL	233
	286	Milden	NIL	342
	314	Dundurn	N/A	173
	344	Corman Park	32	250
	345	Perdue	38	311
	376 A	Eagle Creek	20	252
	376 B	Eagle Creek	15	249
	403	Rosthern	22	252
7A	260 A	Newcombe	N/A	102
	260 B	Newcombe	16.5	211
	261	Chesterfield	N/A	58
	287	St. Andrews	N/A	180
	288	Pleasant Valley	10	159
	290 A	Kindersley	9.1	184
	290 B	Kindersley	N/A	66
	292	Milton	N/A	180
	317 A	Merriott	17	318
	317 B	Marriott	31	301
	318 A	Mountain View	N/A	N/A
	318 B	Mountain View	4	273
	320 A	Oakdale	12	252
	320 B	Oakdale	17	196
	321	Prairiedale	29	254

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	25	255
	347 B	Biggar	58	267
	350 A	Manposa	6	210
	350 B	Manposa	N/A	236
	351	Progress	2	215
	352	Heart's Hill	11	294
	377	Glenside	10	292
	378 A	Rosemount	27	278
	378 B	Rosemount	6	251
	379	Reford	23	239
	380	Tramping Lake	13	175
	381	Grass Lake	N/A	221
	409 A	Buffalo	N/A	232
	409 B	Buffalo	2	236
	410	Round Valley	NIL	283
	411	Senlac	N/A	241
8A	395	Porcupine	N/A	271
	397	Barrier Valley	13.5	289
	428	Star City	7	180
	456	Arborfield	NIL	219
	457	Connaught	11	268
	458	Willow Creek	7	286
	486	Moose Range	NIL	211
	487	Nipawin	3	201
8B	369	St. Peter	2	95
	370	Humboldt	19	164
	371 A	Bayne	7	194
	371 B	Bayne	N/A	N/A
	372	Grant	5.3	269
	402	Fish Creek	10	280
	429	Flett's Springs	8	221
	430	Invergordon	N/A	N/A
	431	St. Louis	N/A	N/A
	459	Kinistino	N/A	224
	460	Birch Hills	N/A	314
9AE	461	Prince Albert	2.5	239
	488 A	Torch River	N/A	316
	488 B	Torch River	N/A	NIL
	491	Buckland	N/A	217
9AW	406	Mayfield	4	211
	435	Redberry	N/A	238
	436	Douglas	NIL	220
	463	Duck Lake	12	239
	464	Leask	14	280
	467 A	Round Hill	N/A	159
	467 B	Round Hill	NIL	235
	494	Canwood	N/A	389
	496	Spiritwood	N/A	53
	497	Medstead	N/A	180
9B	438	Battle River	4	232
	440	Hillsdale	N/A	186
	442	Manitou Lake	0.3	240
	468 A	Meota	N/A	32
	468 B	Meota	N/A	285
	468 C	Meota	5.1	270
	498 A	Parkdale	2	385
	498 B	Parkdale	1	233
	499	Marvin	NIL	322
	501 A	Frenchman Butte	NIL	280
	501 B	Frenchman Butte	NIL	206
	501 C	Frenchman Butte	7	364
	561	Loon Lake	2	296
	588 A	Meadow Lake	NIL	302
	588 B	Meadow Lake	N/A	396
	588 C	Meadow Lake	NIL	451
	588 D	Meadow Lake	32.2	547
	622	Beaver River	N/A	251

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2011
To: August 15, 2011



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2011 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 3.0, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 17, 2011

Cropland Topsoil Moisture Conditions

August 16, 2011



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2011 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 17, 2011

Hay and Pasture Topsoil Moisture Conditions

August 16, 2011



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2011 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 17, 2011